

TOWARDS THE FULL MORDELL-LANG CONJECTURE FOR DRINFELD MODULES

DRAGOS GHIOCA

ABSTRACT. Let ϕ be a Drinfeld module of generic characteristic, and let X be a sufficiently generic affine subvariety of $\mathbb{G}_a^g$. We show that the intersection of X with a finite rank ϕ -submodule of $\mathbb{G}_a^g$ is finite.

1. INTRODUCTION

In [18], McQuillan proved the Mordell-Lang conjecture in its most general form.

Theorem 1.1 (The full Mordell-Lang theorem). *Let G be a semi-abelian variety defined over a number field K . Let $X \subset G$ be a K^{alg} -subvariety, and let $\Gamma \subset G(K^{\text{alg}})$ be a finite rank group (i.e. Γ lies in the divisible hull of a finitely generated subgroup of $G(K^{\text{alg}})$). Then there exist algebraic subgroups $B_1, \dots, B_l$ of G and there exist $\gamma_1, \dots, \gamma_l \in \Gamma$ such that*

$$X(K^{\text{alg}}) \cap \Gamma = \bigcup_{i=1}^l (\gamma_i + B_i(K^{\text{alg}})) \cap \Gamma.$$

We note that in Theorem 1.1 if X does not contain any translate of a positive dimensional algebraic subgroup of G , then the full Mordell-Lang theorem says that $X(K^{\text{alg}}) \cap \Gamma$ is finite. Also, a particular case of the full Mordell-Lang theorem (in the case Γ is the torsion subgroup G_{tor} of G) is the Manin-Mumford theorem, which was first proved by Raynaud [19].

Faltings [8] proved the Mordell-Lang conjecture for finitely generated subgroups Γ of abelian varieties G . His proof was extended by Vojta [23] to finitely generated subgroups of semi-abelian varieties G . Finally, McQuillan [18] extended Vojta's result to finite rank subgroups Γ of semi-abelian varieties G . Later, Rössler [20] provided a simplified proof of McQuillan's extension in which he used uniformities for the intersection of translates of a fixed subvariety $X \subset G$ with the torsion subgroup of the semi-abelian variety G . Essentially, Rössler showed that the full Mordell-Lang conjecture follows from the Mordell-Lang statement for finitely generated subgroups, combined with a uniform Manin-Mumford statement as proved by Hrushovski [17].

It is important to note that the exact translation of the Mordell-Lang conjecture to semi-abelian varieties in characteristic p is false due to the presence of isotrivial varieties. However, Hrushovski [16] saved the Mordell-Lang theorem for finitely generated subgroups of semi-abelian varieties in characteristic p by treating isotrivial varieties as *special*. The isotrivial case was treated by Rahim Moosa and the author in [12] where it was obtained a full Mordell-Lang statement for isotrivial semi-abelian varieties in characteristic p . On the other hand, if we replace G by a power $\mathbb{G}_a^g$ of the additive group scheme, then the exact translation of the Mordell-Lang conjecture either fails (in characteristic 0) or it is trivially true (in characteristic p).

Inspired by the analogy between abelian varieties in characteristic 0 and Drinfeld modules of generic characteristic, Denis [6] proposed that analogs of the Manin-Mumford and Mordell-Lang theorems hold for such Drinfeld modules ϕ acting on $\mathbb{G}_a^g$ (in characteristic p). Denis conjectures ask for describing the intersection of an affine subvariety $X \subset \mathbb{G}_a^g$ with a finite rank ϕ -submodule Γ of $\mathbb{G}_a^g$. Using methods of model theory, combined with some clever number theoretical arguments, Scanlon [21] proved the Denis-Manin-Mumford conjecture. In [9], the author proved the Denis-Mordell-Lang conjecture for finitely generated ϕ -modules Γ under two mild technical assumptions. In this paper, we extend our result from [9] to finite rank ϕ -submodules Γ .

We also note that recently there have been significant progress on establishing additional links between classical diophantine results over number fields and similar statements for Drinfeld modules. The author proved in [10] an equidistribution statement for torsion points of a Drinfeld module, which is similar to the equidistribution statement established by Szpiro-Ullmo-Zhang [22] (which was later extended by Zhang [26] to a full proof of the famous Bogomolov conjecture). Also, Breuer [3] proved a special case of the

André-Oort conjecture for Drinfeld modules, while special cases of this conjecture in the classical case of a number field were proven by Edixhoven-Yafaev [7] and Yafaev [25]. Bosser [2] proved a lower bound for linear forms in logarithms at an infinite place associated to a Drinfeld module (similar to the classical result obtained by Baker [1] for usual logarithms, or by David [4] for elliptic logarithms). Bosser's result was used by Thomas Tucker and the author in [13] to establish certain equidistribution and integrality statements for Drinfeld modules. Moreover, Bosser's result is quite possibly true also for linear forms in logarithms at finite places for a Drinfeld module. Assuming this last statement, Thomas Tucker and the author proved in [14] the analog of Siegel's theorem for finitely generated ϕ -submodules. We believe that our present paper provides an additional proof of the fact that the Drinfeld modules represent the right arithmetic analog in characteristic p for semi-abelian varieties in characteristic 0.

The plan for our paper is as follows: in Section 2 we provide the basic notation for our paper, while in Section 3 we prove our main result (Theorem 3.1).

2. THE MORDELL-LANG THEOREM FOR DRINFELD MODULES

First we note that all subvarieties appearing in this paper are considered to be *closed*. We define next the notion of a Drinfeld module.

Let p be a prime and let q be a power of p . Let C be a projective non-singular curve defined over $\mathbb{F}_q$. Let A be the ring of $\mathbb{F}_q$ -valued functions defined on C , regular away from a fixed closed point $\infty \in C$. Let K be a finitely generated field extension of the fraction field $\text{Frac}(A)$ of A . We let K^{alg} be a fixed algebraic closure of K , and let K^{sep} be the separable closure of K inside K^{alg} .

We define the operator τ as the Frobenius on $\mathbb{F}_q$, extended so that for every $x \in K^{\text{alg}}$, we have $\tau(x) = x^q$. Then for every subfield $L \subset K^{\text{alg}}$, we let $L\{\tau\}$ be the ring of polynomials in τ with coefficients from L (the addition is the usual addition, while the multiplication is given by the usual composition of functions).

Following Goss [15], we call a Drinfeld module of generic characteristic defined over K a morphism $\phi : A \rightarrow K\{\tau\}$ for which the coefficient of τ^0 in ϕ_a is a for every $a \in A$, and there exists $a \in A$ such that $\phi_a \neq a\tau^0$. For the remainder of this paper, unless otherwise stated, $\phi : A \rightarrow K\{\tau\}$ is a Drinfeld module of generic characteristic.

A Drinfeld module $\psi : A \rightarrow K^{\text{alg}}\{\tau\}$ is isomorphic to ϕ (over K^{alg}) if there exists a nonzero $\gamma \in K^{\text{alg}}$ such that for every $a \in A$, we have $\psi_a = \gamma^{-1}\phi_a\gamma$.

For every field extension $K \subset L$, the Drinfeld module ϕ induces an action on $\mathbb{G}_a(L)$ by $a * x := \phi_a(x)$, for each $a \in A$. Let g be a fixed positive integer. We extend diagonally the action of ϕ on $\mathbb{G}_a^g$.

The subgroups of $\mathbb{G}_a^g(K^{\text{alg}})$ invariant under the action of ϕ are called ϕ -submodules. For a ϕ -submodule Γ , its full divisible hull is

$$\Gamma \otimes_A \text{Frac}(A) := \{x \in \mathbb{G}_a^g(K) \mid \text{there exists } 0 \neq a \in A \text{ such that } \phi_a(x) \in \Gamma\}.$$

We define the *rank* of a ϕ -submodule $\Gamma \subset \mathbb{G}_a^g(K^{\text{alg}})$ as $\dim_{\text{Frac}(A)} \Gamma \otimes_A \text{Frac}(A)$.

Definition 2.1. An algebraic ϕ -submodule of $\mathbb{G}_a^g$ is an algebraic subgroup of $\mathbb{G}_a^g$ invariant under ϕ .

Denis proposed in Conjecture 2 of [6] the following problem.

Conjecture 2.2 (The full Denis-Mordell-Lang conjecture). *Let $X \subset \mathbb{G}_a^g$ be an affine variety defined over K^{alg} . Let Γ be a finite rank ϕ -submodule of $\mathbb{G}_a^g(K^{\text{alg}})$. Then there exist algebraic ϕ -submodules $B_1, \dots, B_l$ of $\mathbb{G}_a^g$ and there exist $\gamma_1, \dots, \gamma_l \in \Gamma$ such that*

$$X(K^{\text{alg}}) \cap \Gamma = \bigcup_{i=1}^l (\gamma_i + B_i(K^{\text{alg}})) \cap \Gamma.$$

Before stating our result, we need to introduce the following notion.

Definition 2.3. We call the modular transcendence degree of ϕ the smallest integer $d \geq 1$ such that a Drinfeld module isomorphic to ϕ is defined over a field of transcendence degree d over $\mathbb{F}_q$.

In [9] (see Theorem 4.11), the author proved the following result towards Conjecture 2.2.

Theorem 2.4. *With the above notation, assume in addition that the modular transcendence degree of ϕ is at least 2. Let $X \subset \mathbb{G}_a^g$ be an affine subvariety defined over K^{alg} such that there is no positive dimensional algebraic subgroup of $\mathbb{G}_a^g$ whose translate lies inside X . Let Γ be a finitely generated ϕ -submodule of $\mathbb{G}_a^g(K^{\text{alg}})$. Then $X(K^{\text{alg}}) \cap \Gamma$ is finite.*

In Theorem 3.1 we extend the previous result to all finite rank ϕ -submodules Γ .

Remark 2.5. We have two technical conditions in Theorem 2.4 which we will keep also in our extension from Theorem 3.1. The condition that ϕ has modular transcendence degree at least equal to 2 is a mild technical condition, however necessary due to the methods employed in [9]. The condition that X does not contain any translate of a positive dimensional algebraic subgroup of $\mathbb{G}_a^g$ is satisfied by all sufficiently generic affine subvarieties X .

3. PROOF OF OUR MAIN RESULT

We continue with the notation from Section 2. We define the torsion submodule of ϕ as

$$\phi_{\text{tor}} = \{x \in \mathbb{G}_a^g(K^{\text{alg}}) \mid \text{there exists } a \in A \setminus \{0\} \text{ such that } \phi_a(x) = 0\}.$$

Next we state our main result.

Theorem 3.1. *Let K be a finitely generated field of characteristic p and let g be a positive integer. Let $\phi : A \rightarrow K\{\tau\}$ be a Drinfeld module of generic characteristic. Assume the modular transcendence degree of ϕ is at least 2. Let $X \subset \mathbb{G}_a^g$ be an affine subvariety defined over K^{alg} such that there is no positive dimensional algebraic subgroup of $\mathbb{G}_a^g$ whose translate lies inside X . Let Γ be a finitely generated ϕ -submodule of $\mathbb{G}_a^g(K)$, and let $\Gamma' := \Gamma \otimes_A \text{Frac}(A)$. Then $X(K^{\text{alg}}) \cap \Gamma'$ is finite.*

In our proof of Theorem 3.1 we need a uniform version of Scanlon's result from [21]. He proved the Manin-Mumford theorem for Drinfeld modules in the following form (see his Theorem 1).

Theorem 3.2. *Let $\phi : A \rightarrow K\{\tau\}$ be a Drinfeld module and let $X \subset \mathbb{G}_a^g$ be an affine variety defined over K^{alg} . Then there exist algebraic ϕ -submodules $B_1, \dots, B_l$ of $\mathbb{G}_a^g$ and there exist $\gamma_1, \dots, \gamma_l \in \phi_{\text{tor}}^g$ such that*

$$X(K^{\text{alg}}) \cap \phi_{\text{tor}}^g = \bigcup_{i=1}^l (\gamma_i + B_i(K^{\text{alg}})) \cap \phi_{\text{tor}}^g.$$

In Remark 19 from [21], Scanlon notes that his Manin-Mumford theorem for Drinfeld modules holds uniformly in algebraic families of varieties, i.e. if X varies inside an algebraic family of varieties, then there exists a uniform bound on the degrees of the Zariski closures of $X(K^{\text{alg}}) \cap \phi_{\text{tor}}^g$. In particular, we obtain the following uniform statement for translates of X .

Corollary 3.3. *With the notation for ϕ and X as in Theorem 3.2, assume in addition that X contains no translate of a positive dimensional algebraic subgroup of $\mathbb{G}_a^g$. Then there exists a positive integer N such that for every $x \in \mathbb{G}_a^g(K^{\text{alg}})$, the set $(x + X(K^{\text{alg}})) \cap \phi_{\text{tor}}^g$ has at most N elements.*

Proof. Because X contains no translate of a positive dimensional algebraic subgroup of $\mathbb{G}_a^g$, then for every $x \in \mathbb{G}_a^g(K^{\text{alg}})$, the algebraic ϕ -modules B_i appearing in the intersection $(x + X(K^{\text{alg}})) \cap \phi_{\text{tor}}^g$ are all 0-dimensional. In particular, the set $(x + X(K^{\text{alg}})) \cap \phi_{\text{tor}}^g$ is finite. Thus, using the uniformity obtained by Scanlon for his Manin-Mumford theorem, we conclude that the cardinality of $(x + X(K^{\text{alg}})) \cap \phi_{\text{tor}}^g$ is uniformly bounded by some positive integer N , independent of x . $\square$

We will also use the following fact in the proof of our Theorem 3.1.

Fact 3.4. *Let $\phi : A \rightarrow K\{\tau\}$ be a Drinfeld module. Then for every positive integer d , there exist finitely many torsion points x of ϕ such that $[K(x) : K] \leq d$.*

Proof. If $x \in \phi_{\text{tor}}$, then the canonical height $\hat{h}(x)$ of x (as defined in [5] and [24]) equals 0. Also, as shown in [5], the difference between the canonical height and the usual Weil height is uniformly bounded on K^{alg} . Actually, Denis [5] proves this last statement under the hypothesis that $\text{trdeg}_{\mathbb{F}_q} K = 1$. However, his proof easily generalizes to fields K of arbitrarily finite transcendence degree. For this we need the construction of a coherent good set of valuations on K as done in [11] (see also the similar construction of heights from

[24]). Essentially, a coherent good set U_K of valuations on K is a set of defectless valuations satisfying a product formula on K (for more details, we refer the reader to Sections 2 and 3 of [11]). Then Fact 3.4 follows by noting that there are finitely many points of bounded Weil height and bounded degree over the field K (using Northcott's theorem applied to the global function field K).

Moreover, Corollary 4.22 of [11] provides an effective upper bound on the size of the torsion of ϕ over any finite extension L of K in terms of ϕ and the number of places of L lying above places in U_K of bad reduction for ϕ . Because for each field L such that $[L : K] \leq d$, and for each place $v \in U_K$, there are at most d places w of L lying above v , we conclude that there exists an upper bound for the size of torsion of ϕ over all field extensions of degree at most d over K in terms of ϕ , d and the number of places in U_K of bad reduction for ϕ . $\square$

Proof of Theorem 3.1. Because ϕ has generic characteristic, $\Gamma' \subset \mathbb{G}_a^g(K^{\text{sep}})$. Hence $X(K^{\text{alg}}) \cap \Gamma' = X(K^{\text{sep}}) \cap \Gamma'$.

We note that our theorem is equivalent with showing that if X is a positive dimensional irreducible subvariety of $\mathbb{G}_a^g$ satisfying the hypothesis of Theorem 3.1, then $X(K^{\text{sep}}) \cap \Gamma'$ is not Zariski dense in X . Moreover, at the expense of moding out through the stabilizer $\text{Stab}(X)$ of X (which is a finite group according to the hypothesis of Theorem 3.1), we may assume $\text{Stab}(X)$ is trivial. Note that after moding out through the (finite) group $\text{Stab}(X)$, the variety X is still positive dimensional and irreducible. Finally, at the expense of replacing K by a finite extension, we may assume that X is defined over K , and also $\Gamma \subset \mathbb{G}_a^g(K)$.

Our proof follows the argument from [12] (which in turn was inspired by the argument from [20]). We assume by contradiction that $X(K^{\text{alg}}) \cap \Gamma'$ is Zariski dense in X .

We claim that for every $x \in \Gamma'$ and for every $\sigma \in \text{Gal}(K^{\text{sep}}/K)$, we have $\sigma(x) - x \in \phi_{\text{tor}}^g$. Indeed, because $x \in \Gamma'$, then there exists $0 \neq a \in A$ such that $\phi_a(x) \in \Gamma$. Because $\Gamma \subset \mathbb{G}_a^g(K)$ and ϕ is defined over K , then

$$\phi_a(\sigma(x)) = \sigma(\phi_a(x)) = \phi_a(x).$$

Hence $\phi_a(\sigma(x) - x) = 0$, as desired. Therefore, if $x \in X(K^{\text{sep}})$, then $\sigma(x) - x \in (-x + X) \cap \phi_{\text{tor}}^g$ (we also used that X is defined over K , and so, $\sigma(x) \in X$). Using Corollary 3.3, we obtain an upper bound on the number of conjugates of x , which gives us an upper bound, say N , for $[K(x) : K]$. Implicitly, we also get $[K(\sigma(x)) : K] \leq N$. Because

$$K(\sigma(x) - x) \subset K(x, \sigma(x)),$$

we conclude $[K(\sigma(x) - x) : K] \leq N^2$. Using that $\sigma(x) - x \in \phi_{\text{tor}}^g$, and using Fact 3.4, we conclude that the set

$$(1) \quad \{\sigma(x) - x \mid x \in \Gamma' \text{ and } \sigma \in \text{Gal}(K^{\text{sep}}/K)\} \text{ is finite.}$$

Assuming that $X(K^{\text{sep}}) \cap \Gamma'$ is Zariski dense in X (and also using that X is irreducible), then either $X(K) \cap \Gamma'$ is Zariski dense in X , or $(X(K^{\text{sep}}) \setminus X(K)) \cap \Gamma'$ is Zariski dense in X . If the former statement holds, then X has a Zariski dense intersection with a finitely generated group, as $\Gamma' \cap \mathbb{G}_a^g(K)$ is a finite rank subgroup of the tame module $\mathbb{G}_a^g(K)$ (see Theorem 1 of [24]). By Theorem 2.4, $X(K) \cap (\Gamma' \cap \mathbb{G}_a^g(K))$ is finite and hence it cannot be dense in X (because X is assumed to be positive dimensional). Therefore the latter case of the above dichotomy should occur. Using (1), we conclude that there exists a *nonzero* torsion point $y \in \phi_{\text{tor}}^g$ such that the set

$$\{x \in X(K^{\text{sep}}) \cap \Gamma' \mid \sigma(x) - x = y \text{ for some } \sigma \in \text{Gal}(K^{\text{sep}}/K)\}$$

is Zariski dense in X . Therefore, $y \in \text{Stab}(X)$, contradicting the fact that X has trivial stabilizer. This concludes the proof of Theorem 3.1. $\square$

REFERENCES

- [1] A. Baker, *Transcendental number theory*. Cambridge University Press, London-New York, 1975. x+147 pp.
- [2] V. Bosser, *Minorations de formes linéaires de logarithmes pour les modules de Drinfeld*. (French) [Lower bounds of linear forms in logarithms for Drinfeld modules] J. Number Theory **75** (1999), no. 2, 279–323.
- [3] F. Breuer, *The André-Oort conjecture for products of Drinfeld modular curves*. J. reine angew. Math. **579** (2005), 115–144.
- [4] S. David, *Minorations de formes linéaires de logarithmes elliptiques*. (French) [Lower bounds for linear forms in elliptic logarithms] Mém. Soc. Math. France (N.S.) **62** (1995), iv+143 pp.
- [5] L. Denis, *Hauteurs canoniques et modules de Drinfeld*. (French) [Canonical heights and Drinfeld modules] Math. Ann. **294** (1992), no. 2, 213–223.

- [6] L. Denis, *Diophantine geometry on Drinfeld modules*. The arithmetic of function fields (Columbus, OH, 1991), 285–302, Ohio State Univ. Math. Res. Inst. Publ., 2, de Gruyter, Berlin, 1992.
- [7] B. Edixhoven and A. Yafaev, *Subvarieties of Shimura type*. Ann. of Math. (2) **157** (2003), no. 2, 621–645.
- [8] G. Faltings, *The general case of S. Lang’s conjecture*. Barsotti Symposium in Algebraic Geometry (Abano Terme, 1991), 175–182, Perspect. Math., 15, Academic Press, San Diego, CA, 1994.
- [9] D. Ghioca, *The Mordell-Lang Theorem for Drinfeld modules*, Internat. Math. Res. Notices, **53** (2005), 3273–3307.
- [10] D. Ghioca, *Equidistribution for torsion points of a Drinfeld module*. Math. Ann. **336** (2006), 841–865.
- [11] D. Ghioca, *The Lehmer inequality and the Mordell-Weil theorem for Drinfeld modules*. J. Number Theory **122** (2007), 37–68.
- [12] D. Ghioca and R. Moosa, *Division points on subvarieties of isotrivial semiabelian varieties*. to appear in International Mathematics Research Notices.
- [13] D. Ghioca and T. J. Tucker, *Equidistribution and integrality for Drinfeld modules*. submitted for publication, available online at <http://www.math.mcmaster.ca/~dghioca/papers/papers.html>.
- [14] D. Ghioca and T. J. Tucker, *Siegel’s Theorem for Drinfeld modules*. preprint, available online at <http://www.math.mcmaster.ca/~dghioca/papers/papers.html>.
- [15] D. Goss, *Basic structures of function field arithmetic*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 35. Springer-Verlag, Berlin, 1996.
- [16] E. Hrushovski, *The Mordell-Lang conjecture for function fields*, J. Amer. Math. Soc. **9** (1996), no. 3, 667–690.
- [17] E. Hrushovski, *The Manin-Mumford conjecture and the model theory of difference fields*. Annals of Pure and Applied Logic **112** (2001), 43–115.
- [18] M. McQuillan, *Division points on semi-abelian varieties*. Invent. Math. **120** (1995), no. 1, 143–159.
- [19] M. Raynaud, *Sous-variétés d’une variété abélienne et points de torsion*. (French) [Subvarieties of an abelian variety and torsion points]. Arithmetic and Geometry, Vol. I, 327–352, Prog. Math., **35**, Birkhauser, Boston, MA, 1983.
- [20] D. Rössler, *An afterthought on the generalized Mordell-Lang conjecture*. to appear in the Proceedings of the workshop on Model Theory, Algebraic and Analytic Geometry (Newton Institute, Cambridge, 11–15 July 2005).
- [21] T. Scanlon, *Diophantine geometry of the torsion of a Drinfeld module*. J. Number Theory **97** (2002), no. 1, 10–25.
- [22] L. Szpiro, E. Ullmo and S. Zhang, *Équirépartition des petits points*. (French) [Uniform distribution of small points]. Invent. Math. **127** (1997), no. 2, 337–347.
- [23] P. Vojta, *Integral points on subvarieties of semiabelian varieties. I*. Invent. Math. **126** (1996), no. 1, 133–181.
- [24] J. T.-Y. Wang, *The Mordell-Weil theorems for Drinfeld modules over finitely generated function fields*, Manuscripta math. **106** (2001), 305–314.
- [25] A. Yafaev, *A conjecture of Yves André’s*. Duke Math. J. **132** (2006), no. 3, 393–407.
- [26] S. Zhang, *Equidistribution of small points on abelian varieties*. Ann. of Math. (2) **147** (1998), no. 1, 159–165.

DRAGOS GHIOCA, DEPARTMENT OF MATHEMATICS & STATISTICS, HAMILTON HALL, ROOM 218, MCMASTER UNIVERSITY,
1280 MAIN STREET WEST, HAMILTON, ONTARIO L8S 4K1, CANADA
E-mail address: dghioca@math.mcmaster.ca